

NUCLEAR PHYSICS

Low Temperature Electron Capture Beta-Decay

Folan, L.M., Polytechnic Univ., Applied Mathematics and Physics

Tsifrinovich, V.I., Polytechnic Univ., Applied Mathematics and Physics

Sheverev, V.A., Polytechnic Univ., Applied Mathematics and Physics

The basic idea behind our experiment is that the probability of Electron Capture (EC) Beta Decay depends on the hyperfine state of the decaying isotope.¹⁻³ The total decay rate of a sample of an EC isotope would then depend on the populations of the hyperfine states and hence on temperature. The experiment requires placing radioactive sample crystals inside a scintillation detector, in the bore of a high field magnet, and cooling the apparatus to 20 mK using a dilution refrigerator. X-ray and gamma-ray emissions from the samples are used to monitor the EC process as the temperature and magnetic field are varied. Light from the scintillator is taken to a photomultiplier, at room temperature far from field center, using a fiberoptic bundle. During our experiments at NHMFL the detector was found to operate satisfactorily, with a reference source, at a temperature of 125 mK in a 15 T field. We were unable to cool the detector further, however, because of an unexpectedly large heat leak

associated with the fiber bundle. This was a critical problem as the desired effect is expected to become pronounced below 100 mK. A series of supplementary experiments, performed with the assistance of NHMFL staff member Tim Murphy, has shown that the heat leak can be eliminated by using a fiber bundle containing fewer but larger diameter fibers. Continuation of the experiments is planned for next year.

References:

- 1 Folan, L.M., *et al.*, Phys. Rev. Lett., **74**, 499 (1995).
- 2 Folan, L.M., *et al.*, Trans. Am. Nucl. Soc., **72**, 103 (1995).
- 3 Folan, L.M., *et al.*, J. Appl. Phys., **79**, 5716 (1996).